



*Citation for published version:*

Xu, S, Hu, B, Flower, SE, Jiang, Y-B, Fossey, JS, Deng, W-P & James, TD 2013, 'Colorimetric enantioselective recognition of chiral secondary alcohols *via* hydrogen bonding to a chiral metallocene containing chemosensor', *Chemical Communications*, vol. 49, no. 75, pp. 8314-8316. <https://doi.org/10.1039/c3cc43083a>

*DOI:*

[10.1039/c3cc43083a](https://doi.org/10.1039/c3cc43083a)

*Publication date:*

2013

*Document Version*

Publisher's PDF, also known as Version of record

[Link to publication](#)

*Publisher Rights*

CC BY

This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.

**University of Bath**

**Alternative formats**


If you require this document in an alternative format, please contact:  
[openaccess@bath.ac.uk](mailto:openaccess@bath.ac.uk)

**General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Featuring research from the CASE network in the UK and China. Prof. James, his team and collaborators including Prof. Deng and Dr Fossey describe a chiral metallocene-containing sensor for chiral secondary alcohols.

**Colorimetric enantioselective recognition of chiral secondary alcohols via hydrogen bonding to a chiral metallocene containing chemosensor**

Both catalytically active and inactive diastereoisomers of a chiral metallocene-containing nucleophilic catalyst have been shown to function as sensors for chiral secondary alcohols. Since both diastereoisomers can use hydrogen bonding to recognise chiral secondary alcohols but only one diastereoisomer was previously found to be an active catalyst for secondary alcohol acylation points to divergent recognition mechanisms between the two systems and highlights the utility of investigating catalysis and sensing side by side.

*Two articles in this issue of Chem. Commun. are authored by members of the CASE network, these papers feature as front and back cover articles.*

**As featured in:**



See Wei-Ping Deng and  
Tony D. James *et al.*,  
*Chem. Commun.*, 2013, **49**, 8314.

**RSC Publishing**

**[www.rsc.org/chemcomm](http://www.rsc.org/chemcomm)**

Registered Charity Number 207890

Colorimetric enantioselective recognition of chiral secondary alcohols *via* hydrogen bonding to a chiral metallocene containing chemosensor†Su-Ying Xu,<sup>a</sup> Bin Hu,<sup>b</sup> Stephen E. Flower,<sup>a</sup> Yun-Bao Jiang,<sup>c</sup> John S. Fossey,<sup>bd</sup> Wei-Ping Deng<sup>\*b</sup> and Tony D. James<sup>\*a</sup>Cite this: *Chem. Commun.*, 2013, **49**, 8314Received 25th April 2013,  
Accepted 5th June 2013

DOI: 10.1039/c3cc43083a

www.rsc.org/chemcomm


**An operationally simple colorimetric method for enantioselective detection of chiral secondary alcohols *via* hydrogen bonding interactions using a chiral ferrocene derivative is reported.**

This century has seen an increasing demand for determining the concentration and purity of enantiomers due to the importance of enantiopurity in the pharmaceutical industry.<sup>1,2</sup> Chiral molecular recognition systems have been employed, to assess enantiopurity, which exploit both covalent interactions,<sup>3–8</sup> and non-covalent interactions.<sup>9</sup> Among non-covalent recognition systems reported ionic interactions, hydrogen bonding,<sup>10–12</sup>  $\pi$ - $\pi$  interactions,<sup>13</sup> metal coordination<sup>14–16</sup> and hydrophobic interactions have all been shown to be effective, and these interactions have attracted great interests as they are employed for many applications such as self-assembly<sup>17–19</sup> and molecular recognition.<sup>20–30</sup> The hydrogen bond is an important directional inter- or intra-molecular interaction, which is crucial for controlling molecular conformation and molecular aggregation.<sup>12</sup> In the area of molecular recognition, hydrogen bonding controls the strength of binding between ligands and receptors. For example, in biological systems, the binding between a substrate and an enzyme, as well as cell surface recognition, in great degree, depends on the hydrogen bond interactions.<sup>31</sup>

Steiner outlined the palette of hydrogen bonding patterns available including O-H...N and N-H...O/N interactions.<sup>12</sup> These interactions have been extensively explored in the crystal engineering of supramolecular structures,<sup>32,33</sup> catalytic reactions and molecular recognition.<sup>23,34</sup> Ghosh designed a series of pyridine

derivatives for distinguishing carboxylic acids from non-hydroxyl analogues through hydrogen bonding.<sup>23,34</sup> Shinkai introduced chiral acids as templates to create enantiomerically pure aggregated structures using hydrogen bonding interactions, which have the potential for enantioselective sensing of chiral acids.<sup>20</sup> However, to the best of our knowledge, no one has yet used the nitrogen (N)-hydroxyl (C-OH) hydrogen bonding interaction for enantioselective detection of chiral alcohols. Herein, we report a strategy to enantioselectively detect alcohols through hydrogen bond interactions.

Compounds **1a** and **1b** were previously synthesised and tested as enantioselective catalysts for the kinetic resolution of secondary alcohols (Scheme 1 and S1, ESI†). It was especially noteworthy that whilst diastereoisomer **1b** functioned exquisitely as a catalyst for kinetic resolution by acylation diastereoisomer **1a** was completely inactive (an open top face was reasoned to be required for the acylated catalyst to be able to effectively deliver its cargo). The imidazole nitrogens on **1a** and **1b** are strongly Lewis basic and can themselves form a hydrogen bond with alcohols.<sup>37–39</sup> Therefore, we decided to investigate whether **1a** and **1b** were able to enantioselectively recognise chiral alcohols. From the outset it was observed that the hydrogen bond interactions between **1a** and chiral alcohols are strongly dependent on the solvent, spectral changes in acetonitrile are the most pronounced (S2, ESI†). The binding between **1a** and a series of chiral alcohols was investigated using UV-vis spectroscopy. With dimethyl D/L-tartrates (D-DT and L-DT) the absorption peak was red-shifted from 516 nm to 576 nm (Fig. 1). Enantioselectivity was observed as dimethyl D-tartrate produced



**Scheme 1** Structures of compounds **1a** and **1b**.

<sup>a</sup> Department of Chemistry, University of Bath, Bath BA2 7AY, UK.


E-mail: t.d.james@bath.ac.uk

<sup>b</sup> Shanghai Key Laboratory of Functional Materials Chemistry & School of Pharmacy, East China University of Science and Technology, Shanghai, 20023, China


<sup>c</sup> Department of Chemistry, College of Chemistry and Chemical Engineering, and the MOE Key Laboratory of Analytical Sciences, Xiamen University, Xiamen 361005, China

<sup>d</sup> School of Chemistry, University of Birmingham, Edgbaston, Birmingham B15 2TT, UK

† Electronic supplementary information (ESI) available: Solvent screen and investigations compound **1a**. See DOI: 10.1039/c3cc43083a




**Fig. 1** (a) UV spectra changes of 0.1 mM **1a** in MeCN upon addition of dimethyl D-tartrate; (b) the ratio of absorbance at 576 nm to 516 nm versus concentration of D/L-tartrate for **1a** and **1b**.



**Fig. 2** (a) CD spectra changes of 0.1 mM **1a** in MeCN upon the addition of dimethyl D-tartrate; (b) the ratio of absorbance at 576 nm to 516 nm versus concentration of D/L-tartrate.

larger spectral shifts with **1a** and **1b** than that of dimethyl L-tartrate (Fig. 1 and 2). The observed binding constants for dimethyl D/L-tartrates with **1a** and **1b** are  $392.5 \pm 63.2/112.5 \pm 29.3 \text{ dm}^3 \text{ mol}^{-1}$  and  $298.3 \pm 84.7/141.4 \pm 26.1 \text{ dm}^3 \text{ mol}^{-1}$  respectively (S3, ESI†). Meanwhile enantioselective recognition could be observed colorimetrically, since after the addition of six equivalents of dimethyl D-tartrate to a solution of **1a**, a colour difference is observed as shown in (Fig. 3).

The sensing behaviours of the diastereoisomers **1a** and **1b** with dimethyl D/L-tartrates were identical, demonstrating a divergence between applications to alcohol recognition as opposed to acylation catalysis, *i.e.* the inactive catalyst, **1a**, works equally well as a sensor. As such we chose to make further use of the inactive catalyst and continued our investigations with compound **1a** only. A series of chiral ester containing secondary alcohols were studied and chiral alcohols with  $pK_a \leq 12$  displayed higher chiral discrimination  $\Delta \geq 0.1$  ( $\Delta$  is the difference between  $(A_{576\text{ nm}}/A_{516\text{ nm}})$  for each pair of enantiomers with 0.1 mM **1a** and 0.6 mM of the chiral alcohol)<sup>40</sup> (Table 1). D/L-Tartaric acids were also investigated and while pronounced spectral and colorimetric



**Fig. 3** From left to right: 0.1 mM **1a**, 6 eq. dimethyl L-tartrate, 6 eq. dimethyl D-tartrate in MeCN.

**Table 1** Structures of the chiral secondary alcohols tested in this study and the ratio of absorption at 576 nm to 516 nm of compound **1a** after addition of 6 equivalents of each chiral alcohol

Chiral alcohols ( $A_{576\text{ nm}}/A_{516\text{ nm}}$ ) <sup>a</sup>	$\Delta$ <sup>b</sup>	$pK_a$ <sup>41</sup>
 (-)-Dimethyl D-tartrate 0.982	 (+)-Dimethyl L-tartrate 0.742	0.240 11.44 ± 0.20
 (-)-Diisopropyl D-tartrate 0.641	 (+)-Diisopropyl L-tartrate 0.446	0.195 11.70 ± 0.20
 (2R,3S)-methyl 2,3-dihydroxy-3-phenylpropanoate 0.366	 (2S,3R)-methyl 2,3-dihydroxy-3-phenylpropanoate 0.350	0.016 12.33 ± 0.20
 (+)-Methyl D-lactate 0.626	 (-)-Methyl L-lactate 0.555	0.071 13.07 ± 0.20
 Methyl (R)-(-)-mandelate 0.580	 Methyl (S)-(+)-mandelate 0.457	0.123 12.19 ± 0.20
 (S)-1-phenylpropane-1,3-diol 0.334	 (R)-1-phenylpropane-1,3-diol 0.333	0.001 13.93 ± 0.20
 (1R,2R,3S,5R)-(-)-Pinenediol 0.348	 (1S,2S,3R,5S)-(+)-Pinenediol 0.333	0.015 14.68 ± 0.60

<sup>a</sup> Ratio of absorption at 576 nm to 516 nm, [**1a**] = 0.1 mM, [chiral alcohols] = 0.6 mM. <sup>b</sup>  $\Delta$  is the difference between  $(A_{576\text{ nm}}/A_{516\text{ nm}})$  for each pair of enantiomers.



changes were observed, no enantioselectivity was detected (S4 and S5, ESI†). The observed binding constants for D/L-tartaric acid with **1a** are  $3131 \pm 1157 \text{ dm}^3 \text{ mol}^{-1}$  and  $4636 \pm 1755 \text{ dm}^3 \text{ mol}^{-1}$  respectively (S6, ESI†).  $^1\text{H}$ NMR titrations of **1a** with dimethyl D-tartrate and D-tartaric acid indicate that similar hydrogen bonding species are responsible for the observed spectral changes (S7–S9, ESI†). We believe that enantioselectivity is controlled by steric demands within the hydrogen bonding complexes formed between the guest (acid or alcohol) and compound **1a** (S9, ESI†). Therefore, the increased distance between **1a** and the chiral centres for the hydrogen bonding complexes formed with the tartaric acid (3 bonds) over the alcohols (2 bonds) explains the lack of enantioselectivity observed between the D/L-tartaric acids.

A simple and enantioselective colorimetric sensing strategy for chiral secondary alcohols has been developed based on chiral chemosensor **1**. The hydrogen bond is crucial for the enantioselectivity with more acidic alcohols exhibiting greater enantio-differentiation upon interaction with **1**. Whilst the planar chirality in the sensors was not the overriding chiral recognition motif it is important to note that the metallocene fragment provides a coloured sensor that was vital in establishing a colorimetric assay and the potential to extend this work into the electrochemical arena in the future.

TDJ and SX are grateful for financial support from China Scholarship Council (CSC) and University of Bath Full Fees Scholarship. The Natural Science Foundation of China fellowship for young foreign scientists (No. 21050110426) provided support (JSF and W-PD). Thanks also go to the Catalysis and Sensing for our Environment (CASE) network for facilitating this collaboration. TDJ and JSF thank ECUST for guest professorships. TDJ thanks Xiamen University for a guest professorship.

## Notes and references

- 1 S. C. Stinson, *Chem. Eng. News*, 2000, **78**, 55–78.
- 2 A. Thayer, *Chem. Eng. News*, 2005, **83**, 49–53.
- 3 J. Z. Zhao, T. M. Fyles and T. D. James, *Angew. Chem., Int. Ed.*, 2004, **43**, 3461–3464.
- 4 J. Z. Zhao, M. G. Davidson, M. F. Mahon, G. Kociok-Kohn and T. D. James, *J. Am. Chem. Soc.*, 2004, **126**, 16179–16186.
- 5 X. Zhang, L. N. Chi, S. M. Ji, Y. B. Wu, P. Song, K. L. Han, H. M. Guo, T. D. James and J. Z. Zhao, *J. Am. Chem. Soc.*, 2009, **131**, 17452–17463.
- 6 Y. B. Wu, H. M. Guo, T. D. James and J. Z. Zhao, *J. Org. Chem.*, 2011, **76**, 5685–5695.
- 7 F. Han, L. N. Chi, X. F. Liang, S. M. Ji, S. S. Liu, F. K. Zhou, Y. B. Wu, K. L. Han, J. Z. Zhao and T. D. James, *J. Org. Chem.*, 2009, **74**, 1333–1336.
- 8 T. D. James, K. R. A. S. Sandanayake and S. Shinkai, *Nature*, 1995, **374**, 345–347.
- 9 G. A. Hembury, V. V. Borovkov and Y. Inoue, *Chem. Rev.*, 2008, **108**, 1–73.
- 10 R. G. Desiraju, *The weak hydrogen bond in structural chemistry and biology*, Oxford University Press, 1999.
- 11 M. C. Etter, *Acc. Chem. Res.*, 1990, **23**, 120–126.
- 12 T. Steiner, *Angew. Chem., Int. Ed.*, 2002, **41**, 48–76.
- 13 C. G. Claessens and J. F. Stoddart, *J. Phys. Org. Chem.*, 1997, **10**, 254–272.
- 14 G. R. Whittell, M. D. Hager, U. S. Schubert and I. Manners, *Nat. Mater.*, 2011, **10**, 176–188.
- 15 J. H. Jia, P. Hubberstey, N. R. Champness and M. Schroder, in *Molecular Networks*, ed. M. W. Hosseini, 2009, vol. 132, pp. 135–161.
- 16 B. Moulton and M. J. Zaworotko, *Chem. Rev.*, 2001, **101**, 1629–1658.
- 17 G. M. Whitesides and B. Grzybowski, *Science*, 2002, **295**, 2418–2421.
- 18 J. Seo, J. W. Chung, E. H. Jo and S. Y. Park, *Chem. Commun.*, 2008, 2794–2796.
- 19 M. D. Yilmaz and J. Huskens, *Soft Matter*, 2012, **8**, 11768–11780.
- 20 T. Ishi-i, M. Crego-Calama, P. Timmerman, D. N. Reinhoudt and S. Shinkai, *J. Am. Chem. Soc.*, 2002, **124**, 14631–14641.
- 21 R. Nandhakumar, J. Ryu, H. Park, L. Tang, S. Choi and K. M. Kim, *Tetrahedron*, 2008, **64**, 7704–7708.
- 22 H. Watarai, K. Mitani, N. Morooka and H. Takechi, *Analyst*, 2012, **137**, 3238–3241.
- 23 K. Ghosh, T. Sen and R. Frohlich, *Tetrahedron Lett.*, 2007, **48**, 2935–2938.
- 24 L. Pu, *Acc. Chem. Res.*, 2012, **45**, 150–163.
- 25 L. Cavallo, M. E. Cucciolito, A. De Martino, F. Giordano, I. Orabona and A. Vitagliano, *Chem.-Eur. J.*, 2000, **6**, 1127–1139.
- 26 M. M. Wanderley, C. Wang, C.-D. Wu and W. Lin, *J. Am. Chem. Soc.*, 2012, **134**, 9050–9053.
- 27 Y. Liu, B. Li, T. Wada and Y. Inoue, *Tetrahedron*, 2001, **57**, 7153–7161.
- 28 H. Y. Cun, Y. L. Wang, B. Yang, L. Zhang, S. X. Du, Y. Wang, K. H. Ernst and H. J. Gao, *Langmuir*, 2010, **26**, 3402–3406.
- 29 A. Mravik, Z. Bocskei, K. Simon, F. Elekes and Z. Izsaki, *Chem.-Eur. J.*, 1998, **4**, 1621–1627.
- 30 A. M. Kelly, Y. Perez-Fuertes, J. S. Fossey, S. L. Yeste, S. D. Bull and T. D. James, *Nat. Protoc.*, 2008, **3**, 215–219.
- 31 R. E. Hubbard and M. K. Haider, in *Encyclopedia of Life Sciences (ELS)*, John Wiley & Sons, Ltd., 2010.
- 32 E. Kolomiets, V. Berl and J. M. Lehn, *Chem.-Eur. J.*, 2007, **13**, 5466–5479.
- 33 V. Berl, I. Huc, R. G. Khoury, M. J. Krische and J. M. Lehn, *Nature*, 2000, **407**, 720–723.
- 34 K. Ghosh and S. Adhikari, *Tetrahedron Lett.*, 2006, **47**, 3577–3581.
- 35 B. Hu, M. Meng, Z. Wang, W. T. Du, J. S. Fossey, X. Q. Hu and W. P. Deng, *J. Am. Chem. Soc.*, 2010, **132**, 17041–17044.
- 36 B. Hu, M. Meng, J. S. Fossey, W. Mo, X. Hu and W.-P. Deng, *Chem. Commun.*, 2011, **47**, 10632–10634.
- 37 V. B. Birman, E. W. Uffman, J. Hui, X. M. Li and C. J. Kilbane, *J. Am. Chem. Soc.*, 2004, **126**, 12226–12227.
- 38 M. Wang, Y.-H. Shi, J.-F. Luo, W. Du, X.-X. Shi, J. S. Fossey and W.-P. Deng, *Catal. Sci. Technol.*, 2011, **1**, 100–103.
- 39 M. Wang, Z. Wang, Y.-H. Shi, X.-X. Shi, J. S. Fossey and W.-P. Deng, *Angew. Chem., Int. Ed.*, 2011, **50**, 4897–4900.
- 40 K. N. Kim, K. C. Song, J. H. Noh and S.-K. Chang, *Bull. Korean Chem. Soc.*, 2009, **30**, 197.
- 41 *Calculated using advanced chemistry development (ACD/Labs) software V11.02 ACD/Labs*, 1994–2013.

